

FINITE DIMENSIONAL HOPF ACTIONS ON ALGEBRAIC QUANTIZATIONS

PAVEL ETINGOF AND CHELSEA WALTON

ABSTRACT. Let k be an algebraically closed field of characteristic zero. In joint work with J. Cuadra [CEW1, CEW2], we showed that a semisimple Hopf action on a Weyl algebra over a polynomial algebra $k[z_1, \dots, z_s]$ factors through a group action, and this in fact holds for any finite dimensional Hopf action if $s = 0$. We also generalized these results to finite dimensional Hopf actions on algebras of differential operators. In this work we establish similar results for Hopf actions on other algebraic quantizations of commutative domains. This includes universal enveloping algebras of finite dimensional Lie algebras, spherical symplectic reflection algebras, quantum Hamiltonian reductions of Weyl algebras (in particular, quantized quiver varieties), finite W-algebras and their central reductions, quantum polynomial algebras, twisted homogeneous coordinate rings of abelian varieties, and Sklyanin algebras. The generalization in the last three cases uses a result from algebraic number theory, due to A. Perucca.

1. INTRODUCTION

Throughout the paper, k will denote an algebraically closed field of characteristic zero. In [EW1, Theorem 1.3], we showed that any semisimple Hopf action on a commutative domain over k factors through a group action. Likewise, it was established in our joint work with Juan Cuadra that the same conclusion holds for semisimple Hopf actions on Weyl algebras $\mathbf{A}_n(k[z_1, \dots, z_s])$ [CEW1, Proposition 4.3]. Moreover, we showed that it holds for any (not necessarily semisimple) finite dimensional Hopf action on $\mathbf{A}_n(k)$ [CEW2, Theorem 1.1], and, more generally, on algebras of differential operators of smooth affine varieties [CEW2, Theorem 1.2]. Finally, in [EW3] we extended these results to

2010 *Mathematics Subject Classification.* 16T05, 16S80, 13A35, 16S38.

Key words and phrases. algebraic quantization, filtered deformation, Hopf algebra action, quantum polynomial algebra, Sklyanin algebra, twisted coordinate ring.

certain finite dimensional Hopf actions on deformation quantizations (i.e., formal quantum deformations) of commutative domains. We say that there is *No Finite Quantum Symmetry* in the settings above.

The goal of this paper is establish No Finite Quantum Symmetry results for finite dimensional Hopf actions on other algebraic quantizations of commutative domains, i.e., quantizations whose parameters are elements of k (rather than formal variables). We now summarize our main results for various classes of algebraic quantizations.

1.1. Semisimple Hopf actions on filtered quantizations. Our first main result concerns Hopf actions on *filtered deformations* (or *filtered quantizations*) of commutative domains, that is, on filtered k -algebras B where the associated graded algebra $\text{gr}(B)$ is a commutative finitely generated domain.

Let B be a \mathbb{Z}_+ -filtered algebra over k such that $\text{gr}(B)$ is a commutative finitely generated domain. We will see that for sufficiently large primes p , the algebra B admits a reduction B_p modulo p , which is a domain over $\overline{\mathbb{F}}_p$. Namely, there exists an R -order $B_R \subset B$ over some finitely generated subring $R \subset k$, and $B_p = B_{\psi,p} := B_R \otimes_R \overline{\mathbb{F}}_p$ for a homomorphism $\psi : R \rightarrow \overline{\mathbb{F}}_p$. (For details on R -orders in B see Section 2.1 below).

Recall that a ring A is *PI* if it satisfies a polynomial identity over \mathbb{Z} . By Posner's and Ore's theorems [Pos, Ore] [MR, Theorem 13.6.5 and Corollary 1.14], a domain A is PI if and only if it is an Ore domain and its division ring of fractions $\text{Frac}(A)$ is a central division algebra. In this case, $\text{Frac}(A)$ is a division ring that is dimension d^2 over its center, where d is the *PI degree* of A [MR, Definition 13.6.7].

Definition 1.1. Given B as above, we say that B is an *algebra with PI reductions* if it admits an order B_R such that B_p is PI for sufficiently large p (with any choice of ψ).¹

Theorem 1.2 (Theorem 2.4). *If B is an algebra with PI reductions, then any semisimple Hopf action on B factors through a group action.*

Note that when the Hopf action preserves the filtration of B , Theorem 1.2 (even without the PI reduction assumption) is proved in [EW1,

¹It follows from Lemma 2.1(ii) below that if this condition holds for one pair (R, B_R) , then the condition holds for all such pairs.

Proposition 5.4]; our main achievement here is that we eliminate this requirement.

A basic example of an algebra with PI reductions is the Weyl algebra $B = \mathbf{A}_n(k)$, and, in fact, the proof of Theorem 1.2 is analogous to the proof of [CEW1, Theorem 4.1] which addresses this case. Moreover, a wide range of filtered quantizations (each defined in Subsection 2.2 below) are algebras with PI reductions, resulting in the following corollary.

Corollary 1.3 (Corollary 2.6). *Let B be one of the following filtered k -algebras:*

- (i) *Any filtered quantization B generated in filtered degree one; in particular, the enveloping algebra $U(\mathfrak{g})$ of a finite dimensional Lie algebra \mathfrak{g} , or the algebra $D_\omega(X)$ of twisted differential operators on a smooth affine irreducible variety X ;*
- (ii) *a finite W -algebra or its quotient by a central character;*
- (iii) *a quantum Hamiltonian reduction of a Weyl algebra by a reductive group action; in particular, the coordinate ring of a quantized quiver variety;*
- (iv) *a spherical symplectic reflection algebra; or*
- (v) *the tensor product of any of the algebras above with any commutative finitely generated domain over k .*

Then, any semisimple Hopf action on B factors through a group action.

Other applications of Theorem 1.2 have been investigated recently by Lomp and Pansera [LP]; for instance, they establish No Finite Semisimple Quantum Symmetry on certain iterated differential operator rings.

Remark 1.4. We do not know if a filtered quantization of a finitely generated commutative domain over k must be an algebra with PI reductions (i.e., if the PI reduction assumption is, in fact, vacuous), see the question in [CEW1, Introduction] and [Et2, Question 1.1]. This is of independent interest in noncommutative ring theory.

1.2. Finite dimensional Hopf actions on filtered quantizations. Similarly to [CEW1, Theorem 4.2], Theorem 1.2 and hence Corollary 1.3 hold for Hopf-Galois actions of any (not necessarily semisimple) finite dimensional Hopf algebra (Theorem 3.1). The proof is parallel to the proofs of Theorem 1.2 and [CEW1, Theorem 4.2].

Moreover, it turns out that even without the Hopf-Galois assumption, Theorem 1.2 extends to non-semisimple Hopf actions for a somewhat more restrictive class of quantizations. To see this, let us recall some algebras introduced in [CEW2].

Notation 1.5 ($B, B_{p^m}, C_m, D_{p^m}, Z, Z(m)$). Let B be a quantization with PI reductions, and let B_{p^m} be the reduction of B modulo p^m . Let C_m be the center of B_{p^m} , $\text{Frac}(C_m)$ be its ring of fractions, and $D_{p^m} := B_{p^m} \otimes_{C_m} \text{Frac}(C_m)$. The PI reduction condition implies that D_{p^m} is the full localization (i.e., ring of fractions) of B_{p^m} . These algebras are defined over the truncated Witt ring $W_{m,p}$ of $\overline{\mathbb{F}}_p$ (cf. [CEW2, Subsections 2.1, 2.3, 2.4]). Let Z be the center of the central division algebra D_p . (Here and below, to lighten the notation, we will often suppress dependence on p .) Let Z_m be the center of D_{p^m} , and let $Z(m)$ be its image in D_p under the map $D_{p^m} \twoheadrightarrow D_p$ (so $Z(1) = Z$). It is easy to see that $Z(m) \subset Z(m-1)$.

Definition 1.6. We say that an algebra B with PI reductions is *non-degenerate* if for almost all p one has $\bigcap_{m \geq 1} Z(m) = \overline{\mathbb{F}}_p$.

Theorem 1.7 (Theorem 3.5). *If B is a nondegenerate algebra with PI reductions, then any finite dimensional Hopf action on B factors through a group action (i.e., the condition that H is semisimple in Theorem 1.2 can be dropped).*

The proof of Theorem 1.7 is similar to the proof of [CEW2, Theorem 1.2].

To illustrate when the nondegeneracy condition holds, recall that $\text{gr}(B)$ carries a natural Poisson bracket. Namely, if B is commutative, this bracket is zero; otherwise, if d is the largest integer such that $[F_i B, F_j B] \subset F_{i+j-d} B$, then for $a_0 \in \text{gr}_i(B)$ and $b_0 \in \text{gr}_j(B)$, the Poisson bracket $\{a_0, b_0\}$ is the projection of $[a, b]$ to $\text{gr}_{i+j-d}(B)$, where $a \in F_i B$, $b \in F_j B$ are any lifts of a_0, b_0 respectively. Thus, $\text{gr}(B) = O(X)$, where X is an irreducible Poisson algebraic variety.

The nondegeneracy assumption is satisfied, in particular, when X is a *generically symplectic* Poisson variety, i.e., one having a symplectic dense open subset; see Theorem 3.6. Therefore, Theorem 1.7 holds for many of the examples of Corollary 1.3 — quantum Hamiltonian reductions of Weyl algebras, central reductions of finite W-algebras, spherical symplectic reflection algebras, and tensor products thereof (see Corollary 3.7).

1.3. Quantum polynomial algebras. For our next main result, we consider finite dimensional Hopf actions on *quantum polynomial algebras* (or *quantized coordinate rings of affine n -space*):

$$k_{\mathbf{q}}[x_1, \dots, x_n] := k\langle x_1, \dots, x_n \rangle / (x_i x_j - q_{ij} x_j x_i),$$

where $\mathbf{q} = (q_{ij})$, $q_{ij} \in k^\times$ with $q_{ii} = 1$ and $q_{ij}q_{ji} = 1$. Thus we can view \mathbf{q} as a point of the algebraic torus $(k^\times)^{n(n-1)/2}$ with coordinates q_{ij} for $i < j$.

There are many examples of semisimple Hopf actions on $k_{\mathbf{q}}[x_1, \dots, x_n]$ that do not factor through group actions; the parameters q_{ij} are roots of unity in these examples. See, for instance, [CKWZ, Theorem 0.4], [EW1, Example 5.10], and [KKZ, Examples 7.4–7.6]. Still, we establish the following result.

Let $\langle \mathbf{q} \rangle$ be the subgroup in $(k^\times)^{n(n-1)/2}$ generated by \mathbf{q} , and let $G_{\mathbf{q}}$ be its Zariski closure. Let $G_{\mathbf{q}}^0$ be the connected component of the identity in $G_{\mathbf{q}}$.

Theorem 1.8 (Theorem 4.1). *Let H be a semisimple Hopf algebra of dimension d . If the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to $d!$, then any H -action on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$ factors through a group action.*

If each q_{ij} is a root of unity of order r_{ij} , then $|G_{\mathbf{q}}/G_{\mathbf{q}}^0| = \text{lcm}\{r_{ij}\}_{i < j}$. In particular, if $n = 2$, i.e. if $B = k\langle x, y \rangle / (xy - qyx)$, then the condition on $\mathbf{q} = q \in k^\times$ in Theorem 1.8 is that the order of q is coprime to $d!$ or infinite. On the other hand, the condition on \mathbf{q} in Theorem 1.8 is also satisfied if each q_{ij} is a non-root of unity and the set of the q_{ij} is multiplicatively independent; here, $|G_{\mathbf{q}}/G_{\mathbf{q}}^0| = 1$. See Example 7.3 for a discussion of how to compute $|G_{\mathbf{q}}/G_{\mathbf{q}}^0|$ in general.

One may compare Theorem 1.8 to a similar result, [CWZ, Theorem 4.3], in the case where the Hopf action preserves the grading of $k_{\mathbf{q}}[x_1, \dots, x_n]$. But note that without the degree-preserving assumption, semisimplicity is still needed in Theorem 1.8; see [EW2], [EW3, Example 3.6] for counterexamples for $n = 1, 3$, respectively.

Moreover, Theorem 1.8 is valid for finite dimensional Hopf algebras in the Hopf-Galois case, where we can replace the condition “coprime to $d!$ ” with “coprime to d ” (Proposition 5.1). Also, Theorem 1.8 generalizes straightforwardly (with the same proof) to actions on the quantum torus $k_{\mathbf{q}}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$.

Another generalization of Theorem 1.8 to the non-semisimple case can be made under a nondegeneracy assumption. Recall that \mathbf{q} may be viewed as a skew-symmetric bicharacter on \mathbb{Z}^n with values in k^\times , with $\mathbf{q}(e_i, e_j) = q_{ij}$ for the standard basis $\{e_i\}$. A bicharacter \mathbf{q} is called *nondegenerate* if the character $\mathbf{q}(a, ?) : \mathbb{Z}^n \rightarrow k^\times$ is non-trivial whenever $a \neq 0$. Note that unlike skew-symmetric bilinear forms (which are always degenerate in odd dimensions), a skew-symmetric bicharacter can be nondegenerate for any $n \geq 2$.

Theorem 1.9 (Theorem 5.2). *Let H be a finite dimensional Hopf algebra of dimension d acting on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$. Assume that the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to $d!$, and \mathbf{q} is a nondegenerate bicharacter. Then, the action of H on B factors through a group algebra.*

It is shown in Example 5.5 that Theorems 1.8 and 1.9 fail when hypotheses are removed; these examples involve actions of the non-semisimple 4-dimensional Sweedler Hopf algebra.

1.4. Twisted homogeneous coordinate rings of abelian varieties and Sklyanin algebras. Let X be an abelian variety over k , \mathcal{L} be an ample line bundle on X , and let $\sigma : X \rightarrow X$ be an automorphism given by translation by a point $s \in X$. Then we can define the twisted homogeneous coordinate ring

$$B(X, \sigma, \mathcal{L}) := \bigoplus_{n=0}^{\infty} H^0(X, \bigotimes_{i=0}^{n-1} (\sigma^i)^* \mathcal{L}),$$

with twisted multiplication $f * g := f(\sigma^n)^*(g)$, where f is of degree n ([AV]). It is well-known that $B(X, \sigma, \mathcal{L})$ is a domain, and if $|\sigma| < \infty$, then $B(X, \sigma, \mathcal{L})$ is a PI domain of PI degree $|\sigma|$.

Let G_σ be the Zariski closure of the subgroup $\{s^i\}_{i \in \mathbb{Z}}$, and let G_σ^0 be the connected component of the identity in G_σ .

Theorem 1.10 (Theorem 6.1). *If H is a semisimple Hopf algebra of dimension d , and if the order of G_σ/G_σ^0 is coprime to $d!$, then any H -action on $B(X, \sigma, \mathcal{L})$ factors through a group action.*

In particular, if the subgroup $\{s^i\}_{i \in \mathbb{Z}}$ is Zariski-dense in X , then any semisimple Hopf action on $B(X, \sigma, \mathcal{L})$ factors through a group action. Moreover, if $X =: E$ is an elliptic curve, the condition on σ in Theorem 1.10 is that the order of σ is coprime to $d!$ or infinite.

Lastly, we study semisimple Hopf actions on another class of quantizations: the 3-dimensional Sklyanin algebras $S(a, b, c)$ (Definition 6.3).

To $S(a, b, c)$, one can naturally associate an elliptic curve $E_{abc} \subset \mathbb{P}_k^2$ and an automorphism σ_{abc} given by translation by a point. (See [ATV, Introduction].)

Theorem 1.11. *If H is a semisimple Hopf algebra of dimension d , and if the order of σ_{abc} is coprime to $d!$ or infinite, then any H -action on $S(a, b, c)$ factors through a group action.*

Remark 1.12. We believe that by adapting the techniques in this work, one could establish a version of Theorem 1.11 for semisimple Hopf actions on other elliptic algebras, such as in [Sk1] (or, see [SS]) and in [EG], [Ode], [Ste]. Further, we believe that under an appropriate nondegeneracy condition, there are No Finite Quantum Symmetry results for such elliptic algebras and for twisted homogeneous coordinate rings $B(X, \sigma, \mathcal{L})$ as above; compare to Theorem 1.9.

Our paper is organized as follows. We discuss semisimple Hopf actions on filtered quantizations in Section 2, non-semisimple Hopf actions on filtered quantizations in Section 3, semisimple Hopf actions on quantum polynomial algebras in Section 4, non-semisimple Hopf actions on quantum polynomial algebras in Section 5, and Hopf actions on twisted homogeneous coordinate rings of abelian varieties and Sklyanin algebras in Section 6. The results of Sections 4-6 rely on a number-theoretic result of Antonella Perucca discussed in the Appendix (Section 7).

The notation and terminology of the introduction is used throughout this work, often without mention.

2. SEMISIMPLE HOPF ACTIONS ON FILTERED QUANTIZATIONS

2.1. The result on semisimple Hopf actions on quantizations with PI reductions. In this section, we study actions of semisimple Hopf algebras H on filtered quantizations B . Throughout this section, we let B denote a \mathbb{Z}_+ -filtered algebra over k such that $\text{gr}(B)$ is a commutative finitely generated domain; such B will be referred to as a filtered quantization.

Our goal is to prove Theorem 1.2 (Theorem 2.4 below). This result was established in [CEW1] for B being a Weyl algebra and we generalize the techniques of [CEW1] for our purpose here.

Let R be a finitely generated subring of k . By an R -order in a filtered quantization B we mean an R -subalgebra B_R of B such that $\text{gr}(B_R)$ is a finitely generated R -algebra which is projective as an R -module, and so that the natural map $\text{gr}(B_R) \otimes_R k \rightarrow \text{gr}(B)$ is an isomorphism of graded k -algebras.

Lemma 2.1. (i) *Any filtered quantization B admits an R -order B_R for a suitable ring R .*

(ii) *For any two orders B_R over R and $B_{R'}$ over R' , there exists a finitely generated ring $R'' \subset k$ containing R, R' and an R'' -algebra isomorphism $\phi : B_R \otimes_R R'' \rightarrow B_{R'} \otimes_{R'} R''$ such that $\text{gr}\phi$ is an isomorphism.*

Proof. (i) By the Hilbert basis theorem, the algebra $\text{gr}(B)$ is finitely presented. This implies that so is B , as we can lift the generators and defining relations of $\text{gr}(B)$ to those of B .

More specifically, pick homogeneous generators $\bar{b}_1, \dots, \bar{b}_n$ of $\text{gr}(B)$ of degrees m_1, \dots, m_n . Let

$$p_s(\bar{b}_1, \dots, \bar{b}_n) = 0, \quad s = 1, \dots, r$$

be a set of defining relations for $\text{gr}(B)$, with $p_s \in k[X_1, \dots, X_n]$ being homogeneous of degree d_s (this set may be chosen to be finite by the Hilbert basis theorem). Let b_j be lifts of \bar{b}_j to B , and \tilde{p}_s be homogeneous lifts of p_s to $k\langle X_1, \dots, X_n \rangle$. Then $[b_i, b_j] = f_{ij}(b_1, \dots, b_n)$, where $f_{ij} \in k\langle X_1, \dots, X_n \rangle$ is a noncommutative polynomial of filtration degree at most $m_i + m_j - 1$, and $\tilde{p}_s(b_1, \dots, b_n) = p'_s(b_1, \dots, b_n)$, where $p'_s \in k\langle X_1, \dots, X_n \rangle$ is a noncommutative polynomial of filtration degree at most $d_s - 1$.

Let $g_s := \tilde{p}_s - p'_s \in k\langle X_1, \dots, X_n \rangle$. Thus, we have relations

$$(1) \quad [b_i, b_j] = f_{ij}(b_1, \dots, b_n) \quad \text{and} \quad g_s(b_1, \dots, b_n) = 0$$

in B . It is easy to see that these relations are defining, since they are already defining at the graded level.

Using relations (1), we can find a suitable finitely generated subring $R \subset k$ and define $B_R \subset B$ as follows. We take \tilde{R} to be the ring generated by all the coefficients of the polynomials f_{ij} , g_s , and set $R = \tilde{R}[1/f]$ for a suitable $f \in \tilde{R}$. Now let B_R be the subalgebra of B generated over R by b_1, \dots, b_n . We can choose f so that (1) are defining relations for B_R , and so that B_R is an R -order on B , since for a suitable choice of f , $\text{gr}(B_R)$ is a projective (in fact, free) R -module by

Grothendieck's Generic Freeness Lemma ([Eis, Theorem 14.4]). This proves (i).

(ii) Note that we have a natural isomorphism of filtered algebras $\tilde{\phi} : B_R \otimes_R k \rightarrow B_{R'} \otimes_{R'} k$ (as both are equal to B). This isomorphism is defined over some finitely generated ring $R'' \subset k$ containing R, R' , which implies (ii). \square

Lemma 2.2. *Suppose that B is a filtered quantization that carries an action of a finite dimensional Hopf algebra H . Let S be a finitely generated subring of k , and B_S be an S -order in B . Then, one can find a finitely generated subring $R \subset k$ containing S and a Hopf order $H_R \subset H$ (cf. [CEW1, Definition 2.1]) so that there is an induced action of H_R on $B_R := B_S \otimes_S R$ which gives the action of H on B upon tensoring over R with k .*

Proof. We use the method of [CEW1, Section 2]. Pick homogeneous generators $\bar{b}_1, \dots, \bar{b}_n$ of $\text{gr}(B_S)$, and let b_j be lifts of \bar{b}_j to B_S . Choose a basis $\{h_m\}$ of H . We have

$$(2) \quad h_m \cdot b_j = q_{mj}(b_1, \dots, b_n)$$

for some noncommutative polynomials $q_{mj} \in k\langle X_1, \dots, X_n \rangle$. Let R be generated over S by the structure constants of H in the basis $\{h_m\}$ and the coefficients of q_{mj} . Let $H_R \subset H$ be the span of h_m over R . Then, $H_R \subset H$ is a Hopf order, and H_R acts on B_R by formula (2). The lemma is proved. \square

Thus, any action of H on B admits an R -order for some finitely generated ring $R \subset k$. Moreover, it is easy to see from Lemma 2.1(ii) that any two such orders over rings R, R' can be identified after tensoring with some finitely generated ring $R'' \subset k$ containing R, R' , so an order is essentially unique.

Now fix a ring R and an R -order $B_R \subset B$ with an action of H_R . Let p be a sufficiently large prime, and $\psi : R \rightarrow \overline{\mathbb{F}}_p$ be a character. Following [CEW1, Section 2], let $H_p := H_R \otimes_R \overline{\mathbb{F}}_p$, $B_p := B_R \otimes_R \overline{\mathbb{F}}_p$ be the corresponding reductions of H, B modulo p , where $\overline{\mathbb{F}}_p$ is an R -module via ψ . Then, H_p acts on B_p (by applying ψ to the action of H_R on B_R).

Lemma 2.3. *For a sufficiently large prime p , $\text{gr}(B_p)$, and hence B_p , is a domain.*

Proof. We have $\mathrm{gr}(B_p) = \mathrm{gr}(B_R) \otimes_R \overline{\mathbb{F}}_p$. Therefore, the statement follows from [EGA, 9.7.7(i)] (“geometric irreducibility”). \square

Theorem 2.4. *If B is an algebra with PI reductions, then any semisimple Hopf action on B factors through a group action.*

Proof. We may assume without loss of generality that the action of H on B is inner faithful (otherwise we can pass to an action of a quotient Hopf algebra).

Take $p \gg 0$. Then by [CEW1, Proposition 2.4] (which applies with the same proof in our more general situation), H_p acts inner faithfully on B_p . Moreover, as in [CEW1, Lemma 2.5], H_p is a semisimple cosemisimple Hopf algebra over $\overline{\mathbb{F}}_p$.

Since B is an algebra with PI reductions, by Lemma 2.3, the algebra B_p is a PI domain. Let D_p be the division algebra of quotients of B_p . Then by [Et2, Corollary 3.2(ii)], D_p is a central division algebra of degree p^n for some $n \geq 0$ (which may depend on p). Moreover, H_p acts inner faithfully on D_p by [SV, Theorem 2.2].

Since $\deg D_p = p^n$ is coprime to $(\dim H)!$, [CEW1, Proposition 3.3(ii)] implies that H_p is cocommutative. Thus, H is cocommutative (as in the proof of [CEW1, Theorem 4.1]), and therefore is a group algebra. \square

2.2. Some examples of filtered quantizations. As a consequence, Theorem 2.4 applies to semisimple Hopf actions on many classes of filtered quantizations. Namely, we will consider the following examples.

2.2.1. Twisted differential operators. Let X be a smooth affine irreducible algebraic variety over k , and ω a closed 2-form on X . Then we define the *algebra of twisted differential operators* $D_\omega(X)$ to be the algebra generated by $O(X)$ and elements L_v attached k -linearly to vector fields $v \in \mathrm{Der}O(X)$ on X , with defining relations

$$L_{fv} = fL_v, \quad [L_v, f] = v(f), \quad [L_v, L_w] = L_{[v,w]} + \omega(v, w)$$

for $f \in O(X)$, $v, w \in \mathrm{Der}O(X)$. Then, $D_\omega(X)$ carries a filtration defined by $\deg O(X) = 0$ and $\deg L_v = 1$ for $v \in \mathrm{Der}O(X)$, and $\mathrm{gr}(D_\omega(X)) = O(T^*X)$, the algebra of functions on the symplectic variety T^*X .

The filtered algebra $D_\omega(X)$ depends only on the cohomology class $[\omega]$ of ω , and if $[\omega] = 0$, then $D_\omega(X) = D(X)$, the algebra of usual differential operators on X . For more on twisted differential operators, see e.g. [BB, Section 2].

2.2.2. Quantum Hamiltonian reductions. Let G be a reductive algebraic group over k with Lie algebra \mathfrak{g} , and let $(V, (\cdot, \cdot))$ be a faithful finite dimensional symplectic representation of G . Let $\mathbf{A}(V)$ be the Weyl algebra of V , generated by $v \in V$ with relations $[v, w] = (v, w)$ for $v, w \in V$. We have a natural action of G on $\mathbf{A}(V)$ which preserves its filtration. In this case, we have a natural G -equivariant Lie algebra map $\widehat{\mu} : \mathfrak{g} \rightarrow \mathbf{A}(V)$ called the *quantum moment map*, which quantizes the classical moment map $\mu : V \rightarrow \mathfrak{g}^*$ where $\mu(v)(a) = \frac{1}{2}(v, av)$ for $v \in V, a \in \mathfrak{g}$. Now, given a character $\chi : \mathfrak{g} \rightarrow k$, we can define the algebra

$$B(\chi) := [\mathbf{A}(V)/\mathbf{A}(V)(\widehat{\mu}(a) - \chi(a), a \in \mathfrak{g})]^G,$$

called the *quantum Hamiltonian reduction of $\mathbf{A}(V)$ by G using χ* . It inherits a filtration from the Weyl algebra. See [Et1, Chapter 4] for further details.

Assume that the moment map μ is flat, and that the scheme $\mu^{-1}(0)$ is reduced and irreducible (i.e., $\mu^{-1}(0)$ is a reduced irreducible complete intersection). In this case, $X := \mu^{-1}(0)/G$ is an irreducible generically symplectic Poisson variety, and $B(\chi)$ is a filtered quantization of $O(X)$. See [Lo2, Section 2.3] and [Nak, part 2(i)], and references therein for more details.

An interesting special case of this is when $G = (\prod_{i=1}^r GL(V_i))/k^\times$, and $V = \oplus_{i,j}(V_i \otimes V_j^*)^{\oplus m_{ij}}$, where V_i are finite dimensional vector spaces and $m_{ij} = m_{ji}$ are positive integers where m_{ii} is even; i.e., V is the space of representations of a doubled quiver, and G is the group of linear transformations for this quiver. In this case, $B(\chi)$ is the *quantized quiver variety*, see e.g. [BPW, Section 3.4]. The conditions under which μ is flat and $\mu^{-1}(0)$ is reduced and irreducible are given in [CB, Theorems 1.1 and 1.2].

2.2.3. Finite W -algebras. Let \mathfrak{g} be a simple Lie algebra over k , and $e \in \mathfrak{g}$ be a nilpotent element. To this data one can attach a Lie subalgebra $\mathfrak{m} \subset \mathfrak{g}$ with a character χ , and a *finite W -algebra* is

$$U(\mathfrak{g}, e) := (U(\mathfrak{g})/U(\mathfrak{g})(a - \chi(a), a \in \mathfrak{m}))^{\text{ad } \mathfrak{m}},$$

a quantum Hamiltonian reduction of $U(\mathfrak{g})$. The algebra $U(\mathfrak{g}, e)$ has a filtration induced by the filtration in $U(\mathfrak{g})$, and its associated graded algebra is a polynomial algebra (of functions on the corresponding

Slodowy slice). We refer the reader to [Lo1, Subsections 2.3 and 2.4] and the references therein for details.

Also, the center $U(\mathfrak{g})^{\mathfrak{g}}$ of $U(\mathfrak{g})$ embeds into $U(\mathfrak{g}, e)$, so for any central character $\theta : U(\mathfrak{g})^{\mathfrak{g}} \rightarrow k$ one can consider the central reduction $U_{\theta}(\mathfrak{g}, e) := U(\mathfrak{g}, e)/(a - \theta(a), a \in U(\mathfrak{g})^{\mathfrak{g}})$. Then $\text{gr}(U_{\theta}(\mathfrak{g}, e)) = O(X)$, where X is the nilpotent Slodowy slice, a generically symplectic Poisson variety.

2.2.4. Symplectic reflection algebras. Let G be a finite group and V a faithful finite dimensional symplectic representation of G , and assume that V is not a direct sum of two nonzero symplectic representations. The *symplectic reflection algebra* $H_{t,c}(G, V)$ is the most general filtered deformation of $kG \ltimes SV$, where $[F_i, F_j] \subset F_{i+j-2}$; here $t \in k$ and c is a conjugation invariant function on the set of symplectic reflections in G (see [Et1, Chapter 8]).

Let $\mathbf{e} = |G|^{-1} \sum_{g \in G} g$ be the symmetrizing idempotent for G . Then, the algebra $\mathbf{e}H_{t,c}(G, V)\mathbf{e}$ is called the *spherical symplectic reflection algebra*. For $t = 1$, it is a filtered quantization of $O(X)$ where $X = V/G$, a generically symplectic Poisson variety.

Remark 2.5. There are many other interesting examples of filtered quantizations, and our results given below can be extended to such examples. Since this extension is rather routine, we leave it outside the scope of this paper.

2.3. Results on semisimple Hopf actions on specific filtered quantizations. Here are some concrete applications of Theorem 2.4.

Corollary 2.6. *Let B be one of the following filtered k -algebras:*

- (i) *Any filtered quantization B generated in filtered degree one; in particular, the enveloping algebra $U(\mathfrak{g})$ of a finite dimensional Lie algebra \mathfrak{g} , or the algebra $D_{\omega}(X)$ of twisted differential operators on a smooth affine irreducible variety X ;*
- (ii) *a finite W -algebra or its quotient by a central character;*
- (iii) *a quantum Hamiltonian reduction of a Weyl algebra by a reductive group action; in particular, the coordinate ring of a quantized quiver variety;*
- (iv) *a spherical symplectic reflection algebra; or*
- (v) *the tensor product of any of the algebras above with any commutative finitely generated domain over k .*

Then, any semisimple Hopf action on B factors through a group action.

Note that in some of these cases, a stronger statement is true: any finite dimensional (not necessarily semisimple) Hopf action on B factors through a group action, see Corollary 3.7 below. However, we still prefer to prove the weaker version here, since the proof is simpler (e.g., it does not require reduction modulo prime powers).

Proof. By Theorem 2.4, our job is to show that B is an algebra with PI reductions. In other words, we need to show that the division algebra D_p of fractions of B_p is central (i.e., there is a “ p -center”) for $p \gg 0$. We do so below in each case.

(i) We will show that if a filtered quantization A of a commutative finitely generated domain A_0 over a field F of characteristic $p > 0$ is generated in degree one, then it is module-finite over its center after localization; this implies the required statement.

Let $A_0[i]$ be the degree i part of A_0 . Then $A_0[0] = A[0]$ is a finitely generated commutative domain. Let $\bar{a}_1, \dots, \bar{a}_n$ be generators of $A_0[1]$ as an $A_0[0]$ -module. Let a_i be lifts of \bar{a}_i to A . Then, a_i and $A[0]$ generate A as an algebra. Also the operators $[a_i, ?]$ are derivations of $A[0]$, hence vanish on $A[0]^p$. Thus, $A[0]^p$ is central in A . Let K be the field of quotients of $A[0]^p$, and $A' := A \otimes_{A[0]^p} K$. The K -algebra A' is generated in filtration degree 1, and $L := F_1 A'$ is a finite dimensional vector space over K (as it is spanned by $1, a_1, \dots, a_n$ over $A[0] \otimes_{A[0]^p} K$, and $A[0]$ is module-finite over $A[0]^p$ as $A[0]$ is a finitely generated algebra). Also, L is closed under commutator. Thus, L is a finite dimensional Lie algebra over K , and A' is a quotient of the enveloping algebra $U(L)$. But the enveloping algebra of a finite dimensional Lie algebra in characteristic p is module-finite over its center (i.e., there is a p -center; see [Ja1], [Ja2, Chapter 6, Lemma 5]). This implies that A' is module-finite over its center, as desired.

(ii) Since a W -algebra is a quantum Hamiltonian reduction of the enveloping algebra $U(\mathfrak{g})$ of a semisimple Lie algebra \mathfrak{g} , (see [Lo1]), the statement follows from (i).

(iii) This also follows from (i) and the definition of the quantum Hamiltonian reduction.

(iv) This holds by [BFG, Theorem 9.1.1 (in the appendix)].

(v) This follows easily from the previous cases. □

3. FINITE DIMENSIONAL HOPF ACTIONS ON FILTERED QUANTIZATIONS

3.1. Hopf-Galois actions. Theorem 2.4 does not hold for nonsemisimple Hopf actions, as there are many inner faithful actions of non-semisimple finite dimensional Hopf algebras on commutative domains, see [EW2]. However, Theorem 2.4 is valid in the Hopf-Galois case.

Theorem 3.1. *Let B be a filtered quantization of a commutative finitely generated domain with PI reductions, and let H be a finite dimensional Hopf algebra over k which acts on B . Assume that this action gives rise to an H^* -Hopf-Galois extension $B^H \subset B$. Then, H is a group algebra.*

Proof. The result follows from the arguments in the proofs of Theorem 2.4 and [CEW1, Theorem 4.2]. Namely, recall Notation 1.5. We show similarly to the proof of Theorem 2.4 that Z is H_p -stable, and then proceed as in the proof of [CEW1, Theorem 4.2]. Specifically, by [Et2, Corollary 3.2(ii)], D_p has degree p^n over its center $Z = Z(D_p)$ for some n , so by [CEW1, Proposition 3.3(i)], Z is H_p -invariant. Now, since the action of H on B gives rise to a Hopf-Galois extension, so does the action of H_p on Z , i.e., the algebra map $Z \otimes_{Z^{H_p}} Z \rightarrow Z \otimes H_p^*$ is an isomorphism. Thus, H_p^* is commutative and H_p is cocommutative, so H is cocommutative ([CEW1, Lemma 2.3(ii)], i.e., a group algebra by the Cartier-Gabriel-Kostant theorem [Mo, Corollary 5.6.4(3) and Theorem 5.6.5]. \square

3.2. Preparatory results on nondegenerate quantizations. Another generalization of Theorem 2.4 concerns nondegenerate quantizations, defined in Definition 1.6. To obtain it, we will first need to generalize [CEW2, Theorem 3.2]. Let \mathcal{H} be a finite dimensional Hopf algebra over an algebraically closed field F of characteristic $p > 0$, and let \mathcal{Z} be a finitely generated field extension of F . Let $\mathcal{Z}(m)$, for $m \geq 1$, be a collection of subfields of \mathcal{Z} such that $\mathcal{Z}(m) \supset \mathcal{Z}(m+1)$ for all $m \geq 1$.

Theorem 3.2. *Suppose that $\bigcap_{m \geq 1} \mathcal{Z}(m) = F$, and that $[\mathcal{Z} : \mathcal{Z}(m)]$ is a power of p for all $m \geq 1$. Assume that \mathcal{H} acts F -linearly and inner faithfully on \mathcal{Z} . If $p > \dim \mathcal{H}$ and \mathcal{H} preserves $\mathcal{Z}(m)$ for all m , then \mathcal{H} is a group algebra.*

Proof. The proof is the same as that of [CEW2, Theorem 3.2]. Indeed, the only properties of the fields Z^{p^m} used in that proof are that their intersection is F and that the degree of Z over Z^{p^m} is a power of p . \square

We will also need the lemma below from commutative algebra. Introduce the following notation. Let $W_N = W_N(F) := W(F)/(p^N)$ be the N -th truncated Witt ring of F (W_N is an algebra over $\mathbb{Z}/p^N\mathbb{Z}$; cf. [CEW2, Subsection 2.1]). Let Y be an irreducible smooth affine algebraic variety over F with structure algebra $A := O(Y)$, and \tilde{Y} be a flat deformation of Y over W_N . Let $1 \leq m \leq N$, and let $A_m := O(\tilde{Y})/(p^m)$ (a free $\mathbb{Z}/p^m\mathbb{Z}$ -module); thus $A_1 = A$ and $A_{m-1} = A_m/(p^{m-1})$ for $m \geq 2$. Let

$$d_m : A_m \rightarrow \Omega_{A_m/W_m}$$

be the differential.

Lemma 3.3. *For $1 \leq m \leq N$, the image of $\ker(d_m)$ in A is A^{p^m} .*

Proof. It is clear that the image of $\ker(d_m)$ contains A^{p^m} , so it remains to establish the opposite inclusion. We will do so by induction in m .

The base of induction is the equality $\ker(d|_A) = A^p$, which is the Cartier isomorphism in degree zero ([Kat, Section 7]). Alternatively, here is a direct proof. Since A is integrally closed in its quotient field $L := \text{Frac}(A)$, we may replace A with L . Note that L can be represented as a finite separable extension of $F(y_1, \dots, y_n)$, where $n = \dim Y$. Given $f \in L$ such that $df = 0$, consider the minimal polynomial $P(t) = t^r + a_{r-1}t^{r-1} + \dots + a_0$ of f over $E := F(y_1, \dots, y_n)$. Applying the differential to the equation $P(f) = 0$, we get $\sum_{j=0}^{r-1} f^j da_j = 0$. Since P is the minimal polynomial, this implies that $da_j = 0$ for all j . Thus $a_j \in E^p$ (as the statement in question is easy for purely transcendental fields). Thus, $E^p(f)$ is a finite separable extension of E^p (as P is a separable polynomial). But $E^p(f)$ is a purely inseparable extension of $E^p(f^p)$. Hence, $E^p(f) = E^p(f^p)$, that is, $f \in E^p(f^p) \subset L^p$, as desired.

To perform the induction step, suppose $f \in \ker(d_m)$. Our job is to show that the image \bar{f} of f in A is contained in A^{p^m} . By the induction assumption we know that $\bar{f} = b^{p^{m-1}}$ for some $b \in A$, so it remains to show that $b = c^p$ for some $c \in A$.

For this, let us expand f in a power series in some local coordinate system y_1, \dots, y_n on \tilde{Y} . It is easy to see by looking at monomials that if

$g \in W_m[[y_1, \dots, y_n]]$ and $dg = 0$ then the reduction \bar{g} of g modulo p lies in $F[[y_1^{p^m}, \dots, y_n^{p^m}]]$. In particular, the power series expansion of \bar{f} in y_i is in $F[[y_1^{p^m}, \dots, y_n^{p^m}]]$. This means that the power series expansion of b is in $F[[y_1^p, \dots, y_n^p]]$. Thus, $db = 0$. By the base of induction we conclude that $b = c^p$ for some $c \in A$, which completes the induction step. \square

Moreover, we will need the result below.

Lemma 3.4. *Let B be an algebra with PI reductions, and let D_p denote the full localization (i.e., the ring of fractions) of the reduction B_p of B , for $p \gg 0$. Then, the center Z of D_p is a finitely generated field extension of $\overline{\mathbb{F}}_p$.*

Proof. Let v_1, \dots, v_N be a basis of D_p over Z , and let b_1, \dots, b_n be generators of B_p . Then, $b_s v_i = \sum_{j=1}^N \beta_{si}^j v_j$, for $\beta_{si}^j \in Z$. Let K denote the field $\overline{\mathbb{F}}_p(\beta_{si}^j)$.

Now take $z \in Z$. Then $z \in D_p$, so $z = c^{-1}b$, and hence $cz = b$, for some $b, c \in B_p$ with $c \neq 0$. Since $b, c \in B_p$, they are noncommutative polynomials in b_1, \dots, b_n over $\overline{\mathbb{F}}_p$. So, $bv_i = \sum \beta_i^j v_j$, $cv_i = \sum \gamma_i^j v_j$, with $\beta_i^j, \gamma_i^j \in K$. But $\gamma_i^j z = \beta_i^j$ and γ_i^j are not all zero. So, $z \in K$ and hence $Z = K$. Thus, Z is a finitely generated extension of $\overline{\mathbb{F}}_p$. \square

3.3. Hopf actions on nondegenerate quantizations. Now let B be a filtered quantization with PI reductions.

Theorem 3.5. *If B is a nondegenerate algebra with PI reductions, then any finite dimensional Hopf action on B factors through a group action (i.e., the condition that H is semisimple in Theorem 2.4 can be dropped).*

Proof. The proof is obtained by combining the proofs of Theorem 2.4 and [CEW2, Theorem 1.1] with the following modifications.

1. In [CEW2, Lemma 2.5] and below, x_i, y_i should be replaced by any finite set of generators L_1, \dots, L_r of B , and the number $2n$ in the proof of [CEW2, Lemma 4.3] should be replaced by r (cf. [CEW2, proof of Theorem 1.2]).
2. The discussion in [CEW2, Subsection 2.4, Lemma 4.7, Proposition 4.8] (needed to justify the assumptions of [CEW2, Theorem 3.2]) becomes unnecessary. Instead, note that if $a \in D_{p^m}$ is central modulo p^{m-1} for some $m \geq 2$, then a^p is central. Hence $Z(m) \supset Z(m-1)^p$,

implying that $Z(m) \supset Z^{p^{m-1}}$ and therefore $[Z : Z(m)]$ is finite (by Lemma 3.4) and is a power of p . Now the proof proceeds by invoking Theorem 3.2, whose assumptions are satisfied by the nondegeneracy property of B and using a straightforward generalization of [CEW2, Lemma 4.6].

3. The rest of the proof of [CEW2, Theorem 1.1] is modified as in the proof of Theorem 2.4. Namely, we use the PI reduction condition and [Et2, Corollary 3.2] which says that the PI degree of B_p is a power of p . \square

The next theorem shows that the nondegeneracy assumption is satisfied, in particular, when $\text{gr}(B) = O(X)$, where X is generically symplectic.

Theorem 3.6. *Let B be a quantization with PI reductions, and assume that $\text{gr}(B) = O(X)$, where X is a generically symplectic Poisson variety. Then, any action of a finite dimensional Hopf algebra H on B factors through a group action.*

Proof. By Theorem 3.5, it suffices to show that B is a nondegenerate quantization, i.e., that $\bigcap_{m \geq 1} Z(m) = \overline{\mathbb{F}}_p$, for $p \gg 0$.

Recall Notation 1.5. Let C be the center of B_p ; thus, by Posner's theorem [MR, Theorem 13.6.5], the field $\text{Frac}(C)$ of fractions of C is Z . Let C_m be the center of B_{p^m} , and $C(m)$ be its image in B_p .

Let $a \in B_{p^m}$ be central modulo p (i.e., the image \bar{a} of a in B_p lies in C). Then, a^p is central modulo p^2 , a^{p^2} is central modulo p^3 , and so on. Hence, $C^{p^{m-1}} \subset C(m)$. Let C'_m be the preimage of $C^{p^{m-1}}$ in C_m . Then the image of C'_m in B_p is $C^{p^{m-1}}$.

We claim that

$$(3) \quad Z(m) = \text{Frac}(C(m)).$$

Indeed, it is clear that $\text{Frac}(C(m)) \subseteq Z(m)$. On the other hand, observe that any element $a \in D_{p^m}$ can be written as $a = c^{-1}b$, where $c \in C'_m$ is nonzero modulo p , and $b \in B_{p^m}$ (as this can be done modulo p , since $D_p = Z^{p^{m-1}}B_p$). Now given $z \in Z(m)$, let \tilde{z} be its lift to Z_m . Writing $\tilde{z} = c^{-1}b$ as above, we see that $b := c\tilde{z} \in C_m$. Let $\bar{b} \in C(m)$, $\bar{c} \in C^{p^{m-1}} \subset C(m)$ be the reductions of b, c modulo p , respectively. We have $\bar{b} = \bar{c}\tilde{z}$, hence $z = \bar{c}^{-1}\bar{b} \in \text{Frac}(C(m))$, as claimed.

Now let $B_{0p^m} := \text{gr}(B_{p^m})$. This is a Poisson algebra over the truncated Witt ring $W_{m,p}$. Let C_{0m} be the Poisson center of B_{0p^m} , and

$C_0(m)$ be the image of C_{0m} in B_{0p} . Then $\text{gr}(C_m) \subset C_{0m}$ and hence

$$(4) \quad \text{gr}(C(m)) \subset C_0(m).$$

Let $Z_0 := \text{Frac}(B_{0p})$. Since X is generically symplectic, C_{0m} coincides with the set of all $f \in B_{0p^m}$ such that $df = 0$. By Lemma 3.3 (taking \tilde{Y} to be the reduction modulo p^m of a symplectic dense affine open subset $U \subset X$), this implies that

$$(5) \quad \text{Frac}(C_0(m)) \subset Z_0^{p^m}.$$

Now suppose that $z \in \bigcap_{m \geq 1} Z(m)$ with $z \neq 0$. Then by (3), for each m , we have $z = f_m/g_m$ for $f_m, g_m \in C(m)$. Let $f_m^0, g_m^0 \in \text{gr}(C(m))$ be the leading terms of f_m, g_m . By (4), $f_m^0, g_m^0 \in C_0(m)$. Then for any m, n we have $f_m^0 g_n^0 = f_n^0 g_m^0$ since $f_m g_n = f_n g_m$. So $z_0 := f_m^0/g_m^0$ is independent of m and by (5) belongs to $Z_0^{p^m}$ for all $m \geq 0$. Since $\bigcap_{m \geq 1} Z_0^{p^m}$ is a perfect field that is finitely generated over $\overline{\mathbb{F}}_p$, we get that $\bigcap_{m \geq 1} Z_0^{p^m} = \overline{\mathbb{F}}_p$. So, $z_0 \in \overline{\mathbb{F}}_p$ is a nonzero constant, and $f_m^0 = z_0 g_m^0$ for all m ; in particular,

$$(6) \quad \deg(f_m) = \deg(g_m).$$

Now $z - z_0 = \frac{f_m - z_0 g_m}{g_m}$, and the numerator has degree strictly less than $\deg g_m$. This violates (6), so $z - z_0 = 0$, i.e. $z \in \overline{\mathbb{F}}_p$. This proves the theorem. \square

Corollary 3.7. *Let B be one of the following algebras:*

- (i) *a quotient of a finite W -algebra by a central character;*
- (ii) *a Hamiltonian reduction of a Weyl algebra by a reductive group action; in particular, the coordinate ring of a quantized quiver variety;*
- (iii) *a spherical symplectic reflection algebra $H_{1,c}(G, V)$; or*
- (iv) *the tensor product of any of the algebras in (i)-(iii).*

Then, any action of a finite dimensional Hopf algebra H on B factors through a group action.

Proof. It is explained in Subsection 2.2 that in examples (i)-(iv), we have that $\text{gr}(B) = O(X)$ where X is generically symplectic. This implies the corollary. \square

Proposition 3.8. *Theorems 2.4, 3.1, 3.5, 3.6 remain valid if B is replaced by its quotient division algebra $\text{Frac}(B)$.*

Proof. The proofs are obtained by combining the proofs of Theorem 2.4, 3.1, 3.5, 3.6 with the proof of [CEW1, Proposition 4.4]. (The exact form of the generators of B used in the proof of [CEW1, Proposition 4.4] is irrelevant for the argument.) \square

4. SEMISIMPLE HOPF ACTIONS ON QUANTUM POLYNOMIAL ALGEBRAS

We now turn to finite dimensional Hopf actions on quantum polynomial algebras

$$k_{\mathbf{q}}[x_1, \dots, x_n] := k\langle x_1, \dots, x_n \rangle / (x_i x_j - q_{ij} x_j x_i),$$

where $\mathbf{q} = (q_{ij})$, $q_{ij} \in k^\times$ with $q_{ii} = 1$ and $q_{ij} q_{ji} = 1$. We view \mathbf{q} as a point of the algebraic torus $(k^\times)^{n(n-1)/2}$ with coordinates q_{ij} , $i < j$. Let $\langle \mathbf{q} \rangle$ be the subgroup in $(k^\times)^{n(n-1)/2}$ generated by \mathbf{q} , and let $G_{\mathbf{q}}$ be its Zariski closure. Let $G_{\mathbf{q}}^0$ be the connected component of the identity in $G_{\mathbf{q}}$.

Theorem 4.1. *Let H be a semisimple Hopf algebra of dimension d . If the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to $d!$, then any H -action on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$ factors through a group action.*

Proof. We may assume that H acts on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$ inner faithfully. Let $R \subset k$ be a finitely generated subring containing q_{ij} , let $B_R := R_{\mathbf{q}}[x_1, \dots, x_n]$ be the quantum polynomial algebra defined over R , and H_R be a Hopf R -order with an action on B_R which becomes the action of H on B upon tensoring with k .

Similarly to the proof of Theorem 2.4, we need to control the PI degree of B_R after reducing modulo p ; we employ a version of the number-theoretic result of A. Perucca (as presented in Appendix) to do so.

Given a number field K and a ring homomorphism $\xi : R \rightarrow K$, let $R' := \xi(R)$, $H_{R'} := H_R \otimes_R R'$, $B_{R'} := B_R \otimes_R R' = R'_{\xi(\mathbf{q})}[x_1, \dots, x_n]$. Then, $H_{R'}$ acts on $B_{R'}$ inner faithfully. For a generic choice of ξ , any multiplicative relation satisfied by $\xi(q_{ij})$ is already satisfied by q_{ij} , so by Example 7.3, we have $|G_{\mathbf{q}}/G_{\mathbf{q}}^0| = |G_{\xi(\mathbf{q})}/G_{\xi(\mathbf{q})}^0|$. By Corollary 7.2, there exist infinitely many primes p with prime ideals $\mathfrak{p} \subset R'$ lying over them such that, for a generic homomorphism $\psi : R' \rightarrow \overline{\mathbb{F}}_p$ annihilating \mathfrak{p} , the order $N := N_{\mathfrak{p}}$ of $\psi \circ \xi(\mathbf{q})$ is finite and relatively prime to $d!$. Let $H_p := H_{R'} \otimes_{R'} \overline{\mathbb{F}}_p$ and $B_p := B_{R'} \otimes_{R'} \overline{\mathbb{F}}_p$ be the corresponding

reductions of H and B modulo p . For large enough p , the Hopf algebra H_p is semisimple and cosemisimple (by [CEW1, Lemma 2.5]), and B_p is a PI domain with PI degree dividing N^n (as x_i^N are central elements in B_p). Moreover, H_p acts on B_p inner faithfully by a version of [CEW1, Proposition 2.4] adapted to the algebra B (with the same proof).

Let D_p be the quotient division algebra of B_p . Then, the PI degree of D_p divides N^n , and is therefore coprime to $d!$. Further, H_p acts inner faithfully on D_p . Hence, [CEW1, Proposition 3.3(ii)] implies that H_p is cocommutative. Since this happens for infinitely many primes, we conclude that $H_{R'}$ is cocommutative. Since this happens for generic maps ξ , this implies that H_R is cocommutative. Thus H is cocommutative, i.e., H is a group algebra. \square

Corollary 4.2. *The conclusion of Theorem 4.1 holds when $q_{ij} = q^{m_{ij}}$, where $m_{ij} = -m_{ji}$ are integers, and the order of $q \in k^\times$ is infinite or is coprime to $d!$.*

Proof. This is a special case of Theorem 4.1. \square

Example 4.3. The assumption in Theorem 4.1 and Corollary 4.2 that the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to $d!$ cannot be removed. For instance, there exists an inner faithful action of the 8-dimensional noncommutative noncocommutative semisimple Hopf algebra on the quantum polynomial algebra $k_{-1}[x, y]$, see [KKZ, Example 7.6]. In this case, $|G_{\mathbf{q}}/G_{\mathbf{q}}^0| = 2$.

5. FINITE DIMENSIONAL HOPF ACTIONS ON QUANTUM POLYNOMIAL ALGEBRAS.

Let us now extend the results of the previous section to not necessarily semisimple Hopf algebras, under some additional assumptions.

First of all, when the action of H on B is Hopf-Galois, we can remove in Theorem 4.1 the assumption that H is semisimple, and also weaken the coprimeness assumption, replacing $d!$ with d .

Proposition 5.1. *Suppose that a finite dimensional Hopf algebra H acts on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$, and the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to d . If this action gives rise to an H^* -Hopf-Galois extension $B^H \subset B$, then the action of H on B factors through a group algebra.*

Proof. The proof is parallel to the proof of Theorem 3.1. The weaker coprimeness assumption suffices since by the Hopf-Galois condition,

$[D_p : D_p^H] = d$ (not just $\leq d$). Here, $p \gg 0$ and D_p is the full localization of B reduced modulo p via the method in the proof of Theorem 4.1. \square

Let us now give a generalization of Theorem 4.1 to the non-semisimple case under a nondegeneracy assumption.

Theorem 5.2. *Let H be a finite dimensional Hopf algebra of dimension d acting on $B := k_{\mathbf{q}}[x_1, \dots, x_n]$. Assume that the order of $G_{\mathbf{q}}/G_{\mathbf{q}}^0$ is coprime to $d!$, and \mathbf{q} is nondegenerate. Then, the action of H on B factors through a group action.*

Proof. The proof is obtained by combining the proofs of Theorem 3.5 and Theorem 4.1. Let us describe the necessary changes.

We argue as in the proof of Theorem 4.1. Fix a generic character $\xi : R \rightarrow K$ from R to a number field K , and set $R' = \xi(R)$. By Corollary 7.2, there exist infinitely many primes p with prime ideals $\mathfrak{p} \subset R'$ lying over them such that, for a generic homomorphism $\psi : R' \rightarrow \overline{\mathbb{F}}_p$ annihilating \mathfrak{p} , the order $N := N_{\mathfrak{p}}$ of $\psi \circ \xi(\mathbf{q})$ is finite and coprime to $d!$.

Consider the image $Z(m)$ of the center Z_m of D_{p^m} in D_p (thus, $Z(1) = Z$). By a straightforward generalization of [CEW2, Lemma 4.6], $Z(m)$ is preserved by the action of H_p . It is clear that $Z(m)$ is generated by the monomials $x_1^{m_1} \cdots x_n^{m_n}$ such that $\prod_j q_{ij}^{m_j} = 1$ in the truncated ring of Witt vectors $W_{m,p}$ (see [CEW2, Subsection 2.1]). Let $W'_{m,p}$ be the kernel of the natural map of multiplicative groups $W_{m,p}^{\times} \rightarrow \overline{\mathbb{F}}_p^{\times}$. Then, every element of $W'_{m,p}$ has order a power of p . Hence, $[Z : Z(m)]$ is a power of p . Also it is clear from the nondegeneracy condition for \mathbf{q} that $\bigcap_m Z(m) = \overline{\mathbb{F}}_p$. Thus, Theorem 3.2 applies, and yields that H_p is cocommutative. Hence $H_{R'}$ is cocommutative, implying that H_R is cocommutative and ultimately that H is cocommutative, i.e., a group algebra. \square

Remark 5.3. If $q_{ij} = q^{m_{ij}}$ where q is not a root of unity, then \mathbf{q} is nondegenerate if and only if $\det(m_{ij}) \neq 0$. Theorem 5.2 applies in this case. This gives a generalization of [CWZ, Theorem 0.4] to non-homogeneous Hopf actions for even n .

Proposition 5.4. *Theorem 4.1, Corollary 4.2 and Theorem 5.2 remain valid if the quantum polynomial algebra B is replaced by the quantum torus $k_{\mathbf{q}}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ or the division algebra of quotients $\text{Frac}(B)$.*

Proof. In the case of the quantum torus, the proof is analogous to the proof of Theorem 4.1. The case of the division algebra of quotients is obtained using the same argument as in the proof of [CEW1, Proposition 4.4]. \square

Example 5.5. The condition that H is semisimple cannot be dropped in Theorem 4.1, and the condition that \mathbf{q} is nondegenerate cannot be dropped in Theorem 5.2.

Namely, let $A = A_0 \oplus A_1$ be a $\mathbb{Z}/2\mathbb{Z}$ -graded domain with a nonzero central element $z \in A_1$, and take H to be the 4-dimensional Sweedler Hopf algebra generated by a group-like element g and a $(g, 1)$ -skew-primitive element u with $g^2 = 1$, $u^2 = 0$ and $gu + ug = 0$.

- (1) Then, there is an action of H on A (not preserving the grading of A) given by $g \cdot a = (-1)^{\deg a} a$, and $u \cdot a = 0$ if $a \in A_0$ and $u \cdot a = za$ if $a \in A_1$. It is easy to check that this action is well-defined; it is inner faithful since u acts by a nonzero operator.
- (2) In particular, we have an inner faithful action of H on the quantum polynomial algebra $k_q[x, y]$, for q a root of unity of any odd order $2m - 1$, $m > 0$; namely, we can take $z = x^{2m-1}$.
- (3) Also, this gives an inner faithful action of H on the quantum torus $k_q[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ if n is odd: we can take the central element

$$z = x_1 x_2^{-1} x_3 \cdots x_{n-1}^{-1} x_n.$$

For even n , such an action is impossible if q is not a root of unity by Theorem 5.2. Indeed, the matrix $m_{ij} := \text{sign}(j - i)$ is nondegenerate if and only if n is even (see Remark 5.3).

- (4) Finally, this gives an inner faithful Sweedler Hopf algebra action on the Weyl algebra $\mathbf{A}_n(F)$ when $\text{char}(F) = p \geq 3$; the $\mathbb{Z}/2\mathbb{Z}$ grading is defined by giving the generators degree 1, and we can take, for instance, $z = x_1^p$. (Note that by [CEW2, Theorem 1.1], this is impossible in characteristic zero; indeed, the center of $\mathbf{A}_n(k)$ is k .)

6. SEMISIMPLE HOPF ACTIONS ON TWISTED HOMOGENEOUS COORDINATE RINGS AND 3-DIMENSIONAL SKLYANIN ALGEBRAS

Now let us consider semisimple Hopf actions on twisted homogeneous coordinate rings of abelian varieties. We keep the notation of Subsection 1.4.

Let H be a Hopf algebra over k of dimension d .

Theorem 6.1. *We have the following statements.*

- (i) *If H is semisimple, and if the order of G_σ/G_σ^0 is coprime to $d!$, then any H -action on $B := B(X, \sigma, \mathcal{L})$ factors through a group action.*
- (ii) *Moreover, part (i) holds for the H -action on the division algebra of quotients $\text{Frac}(B)$ of B .*
- (iii) *Part (i) also holds for not necessarily semisimple H if the order of G_σ/G_σ^0 is coprime to d and the H -action gives rise to a Hopf-Galois extension.*

Proof. The proofs of the statements (i),(ii),(iii) are parallel to the proofs of Theorem 4.1, Proposition 5.4, and Proposition 5.1, respectively, where we use that the PI degree of B equals the order of σ . The only difference is that Corollary 7.2 is applied to the abelian variety X with subgroup $\{s^i\}_{i \in \mathbb{Z}}$ rather than the torus $(k^\times)^{n(n-1)/2}$ with subgroup $\langle \mathbf{q} \rangle$. \square

In particular, if $X =: E$ is an elliptic curve, Theorem 6.1 holds if the order of σ is coprime to $d!$ or infinite. Moreover, if σ has infinite order, the assumption that H is semisimple can be dropped.

Theorem 6.2. *Let E be an elliptic curve, and take $\sigma \in \text{Aut}(E)$ given by translation by a point of infinite order. Then, any finite dimensional Hopf action on $B(E, \sigma, \mathcal{L})$ factors through a group action.*

Proof. The proof repeats the proofs of Theorem 5.2 and Theorem 3.5 without significant changes. \square

Finally, let us consider semisimple Hopf actions on three-dimensional Sklyanin algebras [ATV, FO]. Let F be an algebraically closed field of characteristic not equal to 2 or 3.

Definition 6.3. Let $a, b, c \in F^\times$ be such that

$$(3abc)^3 \neq (a^3 + b^3 + c^3)^3.$$

The *three-dimensional Sklyanin algebra*, denoted by $S(a, b, c)$ is generated over F by x, y, z with defining relations

$$ayz + bzy + cx^2 = azx + bxz + cy^2 = axy + byx + cz^2 = 0.$$

It is known that $S(a, b, c)$ is Koszul with Hilbert series $(1 - t)^{-3}$ (see [ATV, Theorems 6.6(ii) and 6.8(i)] and a result of J. Zhang [Smi, Theorem 5.11]), so that $S(a, b, c)$ is a flat deformation of the algebra of polynomials in three variables (see, e.g. [TV, Theorem 1.1]). Moreover, the center of $S(a, b, c)$ contains an element T of degree 3, and $S(a, b, c)/(T) = B(E, \sigma, \mathcal{L})$, where E is the elliptic curve in \mathbb{P}^2 given by the equation

$$(a^3 + b^3 + c^3)xyz = abc(x^3 + y^3 + z^3)$$

with σ given by translation by the point $(a : b : c) \in E$, and \mathcal{L} is a line bundle of degree 3 on E .

Theorem 6.4. *Let $S(a, b, c)$ be a 3-dimensional Sklyanin algebra over k and let H be a semisimple Hopf algebra over k of dimension d . If the order of $\sigma \in \text{Aut}(E)$ is coprime to $d!$ or infinite, then any H -action on $S(a, b, c)$ factors through a group action.*

Proof. It is known from the theory of Sklyanin algebras that if σ has order N then $S(a, b, c)$ is PI with PI degree N (see [AST, part 5 of Theorem on page 7]). Therefore, Theorem 6.4 is proved similarly to Theorem 4.1, using Corollary 7.2 for elliptic curves, as in Theorem 5.2. \square

Remark 6.5. The semisimplicity condition on H in Theorem 6.1 and the infinite order condition in Theorem 6.2 cannot be dropped, as there exists a Sweedler Hopf algebra action on $B := B(X, \sigma, \mathcal{L})$ if σ has odd order N . Namely, we take a sufficiently large odd number m such that the line bundle $\mathcal{L}^{\otimes m}$ is very ample (it exists since \mathcal{L} is ample). Now $B[mN] \neq 0$ and there exists an eigenvector f of σ in $B[mN]$. We then take $z = f^N$, a nonzero central element of odd degree mN^2 , so that a desired action is given by Example 5.5.

Also, the semisimplicity assumption in Theorem 6.4 cannot be dropped, as there exists a Sweedler Hopf algebra action on $S(a, b, c)$ for any a, b, c , given by Example 5.5 where we use the central element T in place of the element z .

7. APPENDIX

The goal of this Appendix is to provide number-theoretic results needed in Section 4. We start with quoting a result from [Per] (in which we take F to be the number field K itself).

Theorem 7.1. [Per, Theorem 7] *Let G be the product of an abelian variety and a torus defined over a number field K . Let $g \in G(K)$ be a K -rational point on G such that the Zariski closure G_g of the subgroup $\langle g \rangle \subset G(K)$ generated by g is connected. Fix a positive integer r . Then there exists a positive Dirichlet density of primes \mathfrak{p} of K such that the order of g modulo \mathfrak{p} is coprime to r . \square*

Corollary 7.2. *Let K, G be as in Theorem 7.1, let $g \in G(K)$, and let $\ell := |G_g/G_g^0|$, where G_g^0 is the connected component of the identity in G_g (i.e., $G_g/G_g^0 = \mathbb{Z}/\ell\mathbb{Z}$). Fix a positive integer r coprime to ℓ . Then, there exists a positive Dirichlet density of primes \mathfrak{p} of K such that the order of g modulo \mathfrak{p} is coprime to r .*

The corollary above is used in the proof of Theorem 4.1 where $d!$ is r and $N_{\mathfrak{p}}$ is the order of g modulo \mathfrak{p} .

Proof. The order of g in G_g/G_g^0 is ℓ , hence $G_{g^\ell} = G_g^0$. Now the statement follows by applying Theorem 7.1 to g^ℓ . \square

Example 7.3. Let G be a split m -dimensional torus, and take an element $g := (q_1, \dots, q_m) \in G$. We have the following statements.

- (1) The group G_g is connected if and only if the group Γ generated by q_1, \dots, q_m in K^\times is free, i.e., does not contain non-trivial roots of unity. Indeed both conditions are equivalent to the condition that any character χ of G which maps g to an ℓ -th root of unity satisfies $\chi(g) = 1$.
- (2) More generally, $|G_g/G_g^0| = \ell$ if and only if the group of roots of unity generated by $\chi(g)$, where χ runs through characters of G such that $\chi(g)$ is a root of unity, is μ_ℓ . In other words, ℓ is the order of the torsion subgroup in \mathbb{Z}^m/g^\perp , where g^\perp is the subgroup of characters χ such that $\chi(g) = 1$. In particular, ℓ depends only on the multiplicative relations satisfied by q_{ij} .
- (3) If $\dim G = 1$ (i.e., $G = \mathbb{G}_m$ or an elliptic curve), then G_g is connected if and only if g has infinite order or $g = 1$. More generally, $|G_g/G_g^0| = \ell > 1$ if and only if g has order ℓ .

ACKNOWLEDGMENTS

We thank Bjorn Poonen for many useful discussions and for the number-theoretic reference [Per], which is crucial for our arguments.

We are also grateful to R. Bezrukavnikov, I. Losev, and H. Nakajima for useful discussions and explanations. We thank the referee for many useful comments that improved greatly the quality of this manuscript. The authors were supported by the National Science Foundation: NSF-grants DMS-1502244 and DMS-1550306.

REFERENCES

- [AST] M. Artin, W. Schelter, and J. Tate, The centers of 3-dimensional Sklyanin algebras, *Barsotti Symposium in Algebraic Geometry: Perspectives in Mathematics*. Vol. 15. Academic Press, 1994.
- [ATV] M. Artin, J. Tate, and M. van den Bergh, Some algebras associated to automorphisms of elliptic curves, *The Grothendieck Festschrift*, Birkhäuser Boston, 1990, 33–85.
- [AV] M. Artin and M. van den Bergh, Twisted homogeneous coordinate rings, *J. Algebra* 133 (1990), 249–271.
- [BB] A. Beilinson and J. Bernstein, A proof of Jantzen conjectures, *Adv. Soviet Math.* 16, Part 1, (1993), 1–50.
- [BFG] R. Bezrukavnikov, M. Finkelberg, and V. Ginzburg (with appendices by P. Etingof and V. Vologodsky), Cherednik algebras and Hilbert schemes in characteristic p , *Represent. Theory* 10 (2006), 254–298.
- [BPW] T. Braden, N. Proudfoot, and B. Webster, Quantizations of conical symplectic resolutions I: local and global structure, arXiv:1208.3863.
- [CKWZ] K. Chan, E. Kirkman, C. Walton, and J. Zhang, Quantum binary polyhedral groups and their actions on quantum planes, to appear in *J. Reine Angew. Math.*, arXiv:1303.7203.
- [CWZ] K. Chan, C. Walton, and J. Zhang, Hopf actions and Nakayama automorphisms, *J. Algebra* 409 (2014), 26–53.
- [CB] W. Crawley-Boevey, Geometry of the moment map for representations of quivers, *Compos. Math.* 126 (2001), no. 3, 257–293.
- [CEW1] J. Cuadra, P. Etingof, and C. Walton, Semisimple Hopf actions on Weyl algebras, *Adv. Math.* 282 (2015), 47–55.
- [CEW2] J. Cuadra, P. Etingof, and C. Walton, Finite dimensional Hopf actions on Weyl algebras, *Adv. Math.* 302 (2016), 25–39.
- [Eis] D. Eisenbud, *Commutative algebra with a view towards algebraic geometry*, Graduate Texts in Mathematics, v.150, 1994.
- [Et1] P. Etingof, *Calogero-Moser systems and representation theory*, European Mathematical Society, 2007.
- [Et2] P. Etingof, A PI degree theorem for quantum deformations, *J. Algebra* 466 (2016), 308–313.
- [EG] P. Etingof and V. Ginzburg, Noncommutative del Pezzo surfaces and Calabi-Yau algebras, *J. Eur. Math. Soc.* 12 (2010), no. 6, 1371–1416.
- [EW1] P. Etingof and C. Walton, Semisimple Hopf actions on commutative domains, *Adv. Math.* 251C (2014), 47–61.

- [EW2] P. Etingof and C. Walton, Pointed Hopf actions on fields, I, *Transform. Groups* 20 (2015), no. 4, 985–1013.
- [EW3] P. Etingof and C. Walton, Finite dimensional Hopf actions on deformation quantizations, arXiv:1602.00532, to appear in *Proc. Amer. Math. Soc.*
- [EGA] A. Grothendieck and J. Dieudonné, Éléments de géométrie algébrique: IV. Étude locale des schémas et des morphismes de schémas, Troisième partie. *Publications Mathématiques de l'IHÉS* 28 (1966), 5–255.
- [FO] B. Feigin and A. Odesskii, Sklyanin elliptic algebras, *Funct. Anal. Appl.* 23 (1989), no. 3, 207–214.
- [Ja1] N. Jacobson, A note on Lie algebras of characteristic p , *Amer. J. Math.* 74 (1952), 357–359.
- [Ja2] N. Jacobson, *Lie algebras*, re-publication of the 1962 original, Dover, New York, 1979.
- [Kat] N. Katz, Nilpotent connections and monodromy theorem, *Publ. Math. IHES* 39 (1970), 175–232.
- [KKZ] E. Kirkman, J. Kuzmanovich, and J. Zhang, Gorenstein subrings of invariants under Hopf algebra actions, *J. Algebra* 322 (2009), no. 10, 3640–3669.
- [LP] C. Lomp and D. Panse, A note on a paper by Cuadra, Etingof, and Walton, arXiv:1506.07766, to appear in *Comm. Alg.*
- [Lo1] I. Losev, Finite W-algebras, *Proc. ICM* (2010), vol. 3, 1281–1307.
- [Lo2] I. Losev, Bernstein inequality and holonomic modules, arXiv:1501.01260.
- [Mo] S. Montgomery, Hopf algebras and their actions on rings, AMS, Providence, 1993.
- [MR] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian rings*, Graduate Studies in Mathematics, 30, American Mathematical Society, Providence, RI, 2001.
- [Nak] H. Nakajima, Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, I, arXiv:1503.03676.
- [Ode] A. Odesskii, Elliptic algebras, (Russian), *Uspekhi Mat. Nauk* 57 (2002), no. 6, 87–122; translation in *Russian Math. Surveys* 57 (2002), no. 6, 1127–1162.
- [Ore] O. Ore, Linear equations in non-commutative fields, *Ann. of Math.* 32 (1931), no. 3, 463–477.
- [Per] A. Perucca, Prescribing valuations of the order of a point in the reductions of abelian varieties and tori, *J. Number Theory* 129 (2009), no. 2, 469–476.
- [Pos] E. Posner, Prime rings satisfying a polynomial identity, *Proc. Amer. Math. Soc.* 11 (1960), 180–183.
- [Sk1] E. Sklyanin, Some algebraic structures connected with the Yang-Baxter equation, *Funct. Anal. Appl.* 16 (1982), no. 4, 263–270.
- [SV] S. Skryabin and F. van Oystaeyen, The Goldie Theorem for H -semiprime algebras, *J. Algebra* 305 (2006), no. 1, 292–320.
- [Smi] S. P. Smith, Some finite-dimensional algebras related to elliptic curves, Representation theory of algebras and related topics (Mexico City, 1994) 19 (1996): 315–348.

- [SS] S. P. Smith and J. T. Stafford, Regularity of the four dimensional Sklyanin algebra, *Compos. Math.* 83 (1992), no. 3, 259–289.
- [Ste] D. Stephenson, Algebras associated to elliptic curves, *Trans. Amer. Math. Soc.* 349 (1997), no. 6, 2317–2340.
- [TV] J. Tate and M. van den Bergh, Homological properties of Sklyanin algebras, *Invent. Math.* 124 (1996), 619–648.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PA 19122, USA

E-mail address: `notlaw@temple.edu`